

Accepted Manuscript


Convolution operators on group algebras which are tauberian or cotauberian

Liliana Cely, Elói M. Galego, Manuel González

PII: S0022-247X(18)30400-1
 DOI: <https://doi.org/10.1016/j.jmaa.2018.05.007>
 Reference: YJMAA 22239

To appear in: *Journal of Mathematical Analysis and Applications*

Received date: 27 November 2017



Please cite this article in press as: L. Cely et al., Convolution operators on group algebras which are tauberian or cotauberian, *J. Math. Anal. Appl.* (2018), <https://doi.org/10.1016/j.jmaa.2018.05.007>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

CONVOLUTION OPERATORS ON GROUP ALGEBRAS WHICH ARE TAUBERIAN OR COTAUBERIAN

LILIANA CELY, ELÓI M. GALEGO, AND MANUEL GONZÁLEZ

ABSTRACT. We study the convolution operators T_μ acting on the group algebras $L_1(G)$ and $M(G)$, where G is a locally compact abelian group and μ is a complex Borel measure on G . We show that a cotauberian convolution operator T_μ acting on $L_1(G)$ is Fredholm of index zero, and that T_μ is tauberian if and only if so is the corresponding convolution operator acting on the algebra of measures $M(G)$, and we give some applications of these results.

1. INTRODUCTION

Tauberian and cotauberian operators were introduced in [14] and [19] respectively, and they have been useful in Banach space theory (see [9]). Recently, some examples of non-trivial (i.e. with non-closed range) tauberian operators $T : L_1(\mu) \rightarrow L_1(\mu)$ have been found in [13], answering a question in [9]. In [3] we considered the case in which T is a convolution operator T_μ acting on the group algebra $L_1(G)$, where G be a locally compact abelian group. We proved that T_μ tauberian implies T_μ invertible when G is non-compact, and that T_μ tauberian implies T_μ Fredholm when G is compact and the singular continuous part μ_{sc} of μ with respect to the Haar measure on G is zero. In particular, tauberian operators T_μ are trivial in those cases. The case $\mu_{sc} \neq 0$ remains open.

Our main result in this paper shows that the cotauberian convolution operators T_μ acting on $L_1(G)$ are always Fredholm, hence trivial. This fact is surprising, because it is not difficult to obtain examples of non-trivial cotauberian operators $T : L_1(\mu) \rightarrow L_1(\mu)$. And it is useful because it gives a new characterization of the Fredholm multipliers of $L_1(G)$ considered in [1, Theorem 5.97] as those that satisfy the definition of cotauberian operator.

In [3], our main tool was a result in [8] characterizing the tauberian operators $T : L_1(\mu) \rightarrow Y$ in terms of the action of T over sequences of normalized disjoint functions in $L_1(\mu)$. Here we follow a more algebraic approach, based in the characterization of the convolution operators T_μ as the multipliers of the group algebra $L_1(G)$ [15, Chapter 0]. We show first that T_μ cotauberian implies T_μ tauberian. In the case G non-compact, T_μ tauberian implies T_μ

Supported in part by MINECO (Spain), Grant MTM2016-76958.
2010 Mathematics Subject Classification. Primary: 47A53, 43A22.
Keywords: multiplier; tauberian operator; convolution operator.

invertible by a result in [3]. In the case G compact, $L_1(G)^{**}$ is a Banach algebra in which $L_1(G)$ is a closed ideal, hence $L_1(G)^{**}/L_1(G)$ is a Banach algebra, and we prove that T_μ cotauberian implies that the operator induced by T_μ^{**} on the quotient algebra $L_1(G)^{**}/L_1(G)$ is bijective. From the inverse of the induced operator we obtain an inverse of T_μ modulo the compact operators, hence T_μ is Fredholm.

We also show that T_μ is tauberian if and only if its natural extension to the algebra of measures $M(G)$ is tauberian, we derive some results for convolution operators acting on $C_0(G)$ and $L_\infty(G)$, and we answer a question raised in [7] about the measures $\mu \in M(G)$ such that $\nu \in M(G)$ and $\mu \star \nu \in L_1(G)$ imply $\nu \in L_1(G)$.

Along the paper X and Y are complex Banach spaces and $T : X \rightarrow Y$ is a (continuous linear) operator from X to Y . We denote by $R(T)$ and $N(T)$ the range and the kernel of T respectively. An operator $T : X \rightarrow Y$ is called *tauberian* if the second conjugate $T^{**} : X^{**} \rightarrow Y^{**}$ satisfies $T^{*-1}(Y) = X$, and T is called *cotauberian* if its conjugate T^* is tauberian.

We denote by X^{co} the quotient space X^{**}/X , and $T^{co} : X^{co} \rightarrow Y^{co}$ is the operator induced by T^{**} and defined by $T^{co}(m + X) := T^{**}m + Y$, $m \in X^{**}$, which is called the *residuum operator* of T . Note that T is tauberian if and only if T^{co} is injective, and T is cotauberian if and only if T^{co} has dense range [9, Proposition 3.1.8 and Corollary 3.1.12]. The set of operators of the form T^{co} for some T was studied in [11].

An operator $T : X \rightarrow Y$ is *Fredholm* if it has closed finite codimensional range and finite dimensional kernel. In this case the *index* of T is defined by $\text{ind}(T) = \dim N(T) - \dim X/R(T)$. Fredholm operators are tauberian and cotauberian [9, Thm. 2.1.5 and Prop. 3.1.5].

2. PRELIMINARIES

We denote by G a locally compact abelian group (a *LCA group*, for short), and by m the Haar measure on G . Moreover $L_1(G)$ is the space of m -integrable complex functions on G endowed with the L_1 -norm $\|\cdot\|_1$, and $M(G)$ denotes the space of complex Borel measures on G endowed with the variation norm. Also $C_0(G)$ and $L_\infty(G)$ denote the space of complex continuous functions on G which vanish at infinity and the space of m -essentially bounded measurable complex valued functions on G , both endowed with the supremum norm. Note that $M(G)$ and $L_\infty(G)$ can be identified with the dual spaces of $C_0(G)$ and $L_1(G)$ respectively, and $L_1(G)$ can be identified with the subspace of those $\mu \in M(G)$ that are absolutely continuous with respect to m .

The space $L_1(G)$ with the convolution $(f \star g)(x) = \int_G f(x-y)g(y)dm(y)$ is a commutative Banach algebra. Moreover, if G is a compact group $L_\infty(G) \subseteq L_1(G)$, thus by Young's inequality, $C(G)$ and $L_\infty(G)$ are commutative Banach algebras with the convolution product. Given $\mu \in M(G)$ and $f \in L_1(G)$, the expression $(\mu \star f)(x) = \int_G f(x-y)d\mu(y)$ defines $\mu \star f \in L_1(G)$

satisfying $\|\mu \star f\|_1 \leq \|\mu\| \cdot \|f\|_1$. Thus we obtain a *convolution operator* T_μ on $L_1(G)$ defined by $T_\mu f = \mu \star f$, and satisfying $\|T_\mu\| = \|\mu\|$. Moreover, given $\mu, \nu \in M(G)$, the convolution of measures $\mu \star \nu \in M(G)$ is commutative [18]. Therefore $T_{\mu \star \nu} = T_\mu T_\nu = T_\nu T_\mu$, and we have an operator $M_\mu : M(G) \rightarrow M(G)$ defined by $M_\mu \lambda = \lambda \star \mu$ which is an extension of T_μ . We can also define the convolution operator $S_\mu : C_0(G) \rightarrow C_0(G)$ and its extension $L_\mu : L_\infty(G) \rightarrow L_\infty(G)$ by $L_\mu g = \mu \star g$.

For $\mu \in M(G)$, we denote by $\tilde{\mu}$ the measure in $M(G)$ such that $\tilde{\mu}(E) = \mu(-E)$ for each Borel set E . Given $\mu, \eta \in M(G)$ and $f \in C_0(G)$, it is not difficult to show that

$$\langle \eta, \mu \star f \rangle = \langle \tilde{\mu} \star \eta, f \rangle.$$

Thus $S_\mu^* = M_{\tilde{\mu}}$, and similarly we get $T_\mu^* = L_{\tilde{\mu}}$.

Following [15], we say that a Banach algebra A is *without order* if for all $x \in A$, $xA = \{0\}$ implies $x = 0$, or, for all $x \in A$, $Ax = \{0\}$ implies $x = 0$. A map $T : A \rightarrow A$ is a *multiplier of A* if $x(Ty) = (Tx)y$ for all $x, y \in A$. If G is a LCA group, then $L_1(G)$ is without order and the multipliers of $L_1(G)$ are precisely the convolution operators T_μ , $\mu \in M(G)$ [15, Chapter 0].

A net $(e_\alpha)_{\alpha \in I}$ in a commutative Banach algebra A is an *approximate identity* if, for each $a \in A$, $\lim_\alpha \|ae_\alpha - a\| = 0$. If G is a LCA group, then $L_1(G)$ admits a bounded approximate identity [15, Appendix F].

Given a Banach algebra A , the second dual space A^{**} of A is also a Banach algebra endowed with the (first) Arens product [2]. Specifically, given $M, N \in A^{**}$, $f \in A^*$ and $a, b \in A$, we define the product $M \cdot N$ in three steps as follows:

$$\begin{aligned} f \cdot a \in A^* : & \quad \langle f \cdot a, b \rangle := \langle f, ab \rangle \\ N \cdot f \in A^* : & \quad \langle N \cdot f, a \rangle := \langle N, f \cdot a \rangle \\ M \cdot N \in A^{**} : & \quad \langle M \cdot N, f \rangle := \langle M, N \cdot f \rangle. \end{aligned}$$

Thus, for G a LCA group, $L_1(G)^{**}$ is a Banach algebra. Given $f \in L_\infty(G)$ and $\phi \in L_1(G)$, it is not difficult to show that

$$(1) \quad f \cdot \phi = f \star \tilde{\phi},$$

where $\tilde{\phi}(x) = \phi(-x)$. Moreover, since G is commutative, the centre of $L_1(G)^{**}$ is $L_1(G)$ [16, Corollary 3] i.e.,

$$L_1(G) = \{m \in L_1(G)^{**} : m \cdot n = n \cdot m \text{ for each } n \in L_1(G)^{**}\}.$$

When G is compact, $L_1(G)$ is a (closed) ideal of $L_1(G)^{**}$ [20, Proposition 4.2]. Thus $L_1(G)^{co}$ is a Banach algebra.

We will need later the following result proved in [3].

Theorem 2.1. [3, Theorem 3.2] *Let G be a non-compact LCA group. Then every tauberian convolution operator $T_\mu : L_1(G) \rightarrow L_1(G)$ is invertible.*

For basic results on Fredholm theory, tauberian operators and multipliers of Banach algebras without order we refer to [1], [9] and [15].

3. MAIN RESULTS

First we show that the Banach algebras involved in our arguments are without order.

Proposition 3.1. *Let G be a LCA group. Then the algebra $L_1(G)^{**}$ admits a norm-one right identity; hence it is a Banach algebra without order. Moreover, when G is compact, the quotient algebra $L_1(G)^{co}$ also admits a norm-one right identity and it is a Banach algebra without order.*

Proof. The first part of our first result was proved in [6, Proposition 2.1]. For the second part, note that if E is a norm-one right identity of $L_1(G)^{**}$ and G is compact, then $L_1(G)^{co}$ is a Banach algebra and $E + L_1(G)$ is a norm-one right identity in $L_1(G)^{co}$, hence $L_1(G)^{co}$ is a Banach algebra without order. \square

Let us see that the multipliers of algebras without order have a good behavior under duality.

Proposition 3.2. *Let T be a multiplier of a Banach algebra without order A . Then the second conjugate $T^{**} : A^{**} \rightarrow A^{**}$ is a multiplier of A^{**} .*

Proof. The proof has three steps. Let $M, N \in A^{**}$, $f \in A^*$ and $a, b \in A$. Then $T(ab) = (Ta)b = a(Tb)$, because A is without order, and from this equality we derive $T^*(f \cdot a) = f \cdot Ta$ as follows:

$$\begin{aligned} \langle T^*(f \cdot a), b \rangle &= \langle f \cdot a, Tb \rangle \\ &= \langle f, a(Tb) \rangle \\ &= \langle f, (Ta)b \rangle \\ &= \langle f \cdot Ta, b \rangle. \end{aligned}$$

In a similar way, from $T^*(f \cdot a) = f \cdot Ta$ we derive $T^{**}N \cdot f = T^*(N \cdot f)$, and from the latter equality we get that $M \cdot T^{**}N = T^{**}M \cdot N$, showing that T^{**} is a multiplier. \square

Corollary 3.3. *Let G be a compact LCA group. Then the residuum operator T_μ^{co} is a multiplier of $L_1(G)^{co}$.*

Proof. Let $n + L_1(G)$ and $m + L_1(G)$ in $L_1(G)^{co}$. Since T_μ^{**} is a multiplier of $L_1(G)^{**}$,

$$\begin{aligned} (n + L_1(G)) \cdot T_\mu^{co}(m + L_1(G)) &= (n + L_1(G)) \cdot (T_\mu^{**}m + L_1(G)) \\ &= n \cdot T_\mu^{**}m + L_1(G) \\ &= T_\mu^{**}n \cdot m + L_1(G) \\ &= (T_\mu^{**}n + L_1(G)) \cdot (m + L_1(G)) \\ &= T_\mu^{co}(n + L_1(G)) \cdot (m + L_1(G)). \end{aligned}$$

Hence T_μ^{co} is a multiplier of $L_1(G)^{co}$. \square

Now we are ready to show that cotauberian convolution operators on $L_1(G)$ are tauberian. This result contrasts with the fact that it is easy to find non-trivial cotauberian operators $T : L_1(0, 1) \rightarrow L_1(0, 1)$: just consider a surjective operator S with infinite dimensional kernel from $L_1(0, 1)$ onto a closed infinite dimensional subspace M of $L_1(0, 1)$ such that the quotient $L_1(0, 1)/M$ is reflexive, and add a compact operator K so that $T = S + K$ has non-closed range. However it is difficult to obtain a non-trivial tauberian operator $T : L_1(0, 1) \rightarrow L_1(0, 1)$ (see [13]).

Proposition 3.4. *Let T be a multiplier of a Banach algebra without order A . Suppose that A coincides with the centre of A^{**} and T is cotauberian. Then T is tauberian.*

Proof. Let $M \in A^{**}$ such that $T^{**}M \in A$. Then for each $K \in A^{**}$,

$$M \cdot T^{**}K = T^{**}M \cdot K = K \cdot T^{**}M = T^{**}K \cdot M$$

because T^{**} is a multiplier of A^{**} . In particular $M \cdot T^{**}K = T^{**}K \cdot M$. Since T is cotauberian, $R(T^{**}) + A$ is dense in A^{**} [9, Corollary 3.1.12]. Consequently, given $S \in A^{**}$, we can find sequences $(W_n) \subset A^{**}$ and $(f_n) \subset A$ with $(T^{**}W_n + f_n)$ norm-convergent to S . Therefore

$$\begin{aligned} M \cdot S &= \lim_n (M \cdot (T^{**}W_n + f_n)) \\ &= \lim_n (M \cdot T^{**}W_n + M \cdot f_n) \\ &= \lim_n (T^{**}W_n \cdot M + f_n \cdot M) \\ &= \lim_n ((T^{**}W_n + f_n) \cdot M) \\ &= S \cdot M. \end{aligned}$$

Then $M \in A$, the centre of A^{**} . Hence T is tauberian. \square

Corollary 3.5. *Let G be a LCA group.*

- (1) *Every cotauberian convolution operator T_μ is tauberian.*
- (2) *Suppose that G is non-compact. Then T_μ is cotauberian if and only if it is invertible.*

Proof. (1) It follows from the facts that $L_1(G)$ is a Banach algebra without order and $L_1(G)$ is the centre of $L_1(G)^{**}$.

(2) It is enough to observe that, since G is non-compact, every tauberian convolution operator on $L_1(G)$ is invertible (Theorem 2.1). \square

Let E be a right identity in the algebra $L_1(G)^{**}$ provided by Proposition 3.1. We consider the map $\Gamma_E : M(G) \rightarrow L_1(G)^{**}$ defined by

$$\Gamma_E(\mu) := T_\mu^{**}(E), \quad \mu \in M(G).$$

The map Γ_E is an isometric algebra homomorphism of $M(G)$ into $L_1(G)^{**}$ extending the natural embedding of $L_1(G)$ into $L_1(G)^{**}$ [6, Proposition 2.3].

Since T_μ^{**} is a multiplier of $L_1(G)^{**}$, for each $m \in L_1(G)^{**}$ we have

$$(2) \quad T_\mu^{**}m = (T_\mu^{**}m) \cdot E = m \cdot T_\mu^{**}E = m \cdot \Gamma_E(\mu).$$

Thus T_μ^{**} is a right multiplication operator (by $\Gamma_E(\mu)$). Moreover

$$(3) \quad E \cdot \Gamma_E(\mu) = T_\mu^{**}(E) = \Gamma_E(\mu).$$

Next we give our main result.

Theorem 3.6. *Let G be a LCA group and let $T_\mu : L_1(G) \rightarrow L_1(G)$ be a convolution operator. Then T_μ is cotauberian if and only if it is Fredholm of index zero.*

Proof. For G non-compact, it is a consequence of Corollary 3.5.

Suppose that G is compact and T_μ cotauberian. Then T_μ is also tauberian by Proposition 3.4, hence T_μ^{co} is injective with dense range by [9, Prop. 3.1.8 and Cor. 3.1.12], and has closed range by [9, Theorem 4.4.2]. Thus T_μ^{co} is invertible.

Since T_μ^{co} is a multiplier of the Banach algebra without order $L_1(G)^{co}$ (Proposition 3.1), its inverse $S := (T_\mu^{co})^{-1}$ is a multiplier of $L_1(G)^{co}$ [15, Theorem 1.1.3].

Let E be a right identity in $L_1(G)^{**}$. Then

$$T_\mu^{co}(E + L_1(G)) = T_\mu^{**}E + L_1(G) = \Gamma_E(\mu) + L_1(G).$$

Set $\beta + L_1(G) := S(E + L_1(G))$. Then

$$\begin{aligned} (\beta + L_1(G)) \cdot (\Gamma_E(\mu) + L_1(G)) &= S(E + L_1(G)) \cdot T_\mu^{co}(E + L_1(G)) \\ &= T_\mu^{co}(S(E + L_1(G))) \cdot (E + L_1(G)) \\ &= E + L_1(G). \end{aligned}$$

Thus $(\beta \cdot \Gamma_E(\mu) - E) \in L_1(G)$. On the other hand, we have

$$\begin{aligned} (E + L_1(G)) \cdot (\beta + L_1(G)) &= (E + L_1(G)) \cdot S(E + L_1(G)) \\ &= S(E + L_1(G)) \cdot (E + L_1(G)) = \beta + L_1(G), \end{aligned}$$

hence $(E \cdot \beta - \beta) \in L_1(G)$. Since $L_1(G)$ is the centre of $L_1(G)^{**}$, for each $n \in L_1(G)^{**}$ we have

$$(E \cdot \beta - \beta) \cdot n = n \cdot (E \cdot \beta - \beta) = n \cdot \beta - n \cdot \beta = 0.$$

Then $E \cdot \beta - \beta = 0$, because $L_1(G)^{**}$ is a Banach algebra without order, thus $E \cdot \beta = \beta$.

Let $\nu \in M(G)$ denote the restriction of $\beta \in L_\infty(G)^*$ to $C(G)$. By [6, Proposition 2.5] we have that $\beta - \Gamma_E(\nu)$ is zero on $C(G)$. Moreover, by formula (3), we have

$$E \cdot (\beta - \Gamma_E(\nu)) = \beta - \Gamma_E(\nu).$$

The right identity $E \in L_1(G)^{**}$ is the w^* -limit of $(i(e_\gamma))$ for some bounded approximate identity $(e_\gamma) \subset L_1(G)$, where $i : L_1(G) \rightarrow L_1(G)^{**}$ is the

canonical inclusion. Therefore, for $f \in L_\infty(G)$ we have

$$\begin{aligned} \langle \beta - \Gamma_E(\nu), f \rangle &= \langle E \cdot (\beta - \Gamma_E(\nu)), f \rangle \\ &= \langle E, (\beta - \Gamma_E(\nu)) \cdot f \rangle \\ &= \lim_{\gamma} \langle i(e_\gamma), (\beta - \Gamma_E(\nu)) \cdot f \rangle \\ &= \lim_{\gamma} \langle (\beta - \Gamma_E(\nu)) \cdot f, e_\gamma \rangle \\ &= \lim_{\gamma} \langle (\beta - \Gamma_E(\nu)), f \cdot e_\gamma \rangle = 0 \end{aligned}$$

because, by (1) and [18, Theorem 1.1.6], $f \cdot e_\gamma \in C(G)$, thus $\beta = \Gamma_E(\nu)$.

Note that $I = T_{\delta_0}$, where I is the identity operator on $L_1(G)$ and δ_0 is the unit measure concentrated at $\{0\}$. Then $T_{\delta_0}^{**}$ is the identity operator on $L_1(G)^{**}$, and thus $E = \Gamma_E(\delta_0)$. Hence, since $\Gamma_E : M(G) \rightarrow L_1(G)^{**}$ is an algebra homomorphism,

$$(4) \quad \Gamma_E(\nu \star \mu - \delta_0) = \Gamma_E(\nu) \cdot \Gamma_E(\mu) - \Gamma_E(\delta_0) = \beta \cdot \Gamma_E(\mu) - E \in L_1(G).$$

By formula (4) and [6, proposition 2.3], $\mu \star \nu - \delta_0 \in L_1(G)$, and thus

$$T_{\mu \star \nu - \delta_0} = T_\mu T_\nu - I$$

is a compact operator [4, Lemma 2.1.4]. Thus $T_\mu T_\nu = T_\nu T_\mu$ is a Fredholm operator, hence T_μ is Fredholm, and it has index zero by [1, Theorem 5.97].

For the converse implication, note that Fredholm operators are trivially cotauberian. \square

Next we study the convolution operator $M_\mu : M(G) \rightarrow M(G)$ and its relation with $T_\mu : L_1(G) \rightarrow L_1(G)$.

Theorem 3.7. *Let G be a LCA group and let $\mu \in M(G)$. Then*

- (1) M_μ cotauberian implies M_μ tauberian.
- (2) T_μ is tauberian if and only if M_μ is tauberian.

Proof. (1) It was shown in [17, Main theorem] that, as in the case of $L_1(G)$, the measure algebra $M(G)$ is the centre of $M(G)^{**}$. Since $M(G)$ has an identity, it is without order. Thus the result follows from Proposition 3.4.

(2) Suppose that T_μ is tauberian, and let E be a right identity in $L_1(G)^{**}$. Then the following diagram is commutative:

$$\begin{array}{ccc} L_1(G)^{**} & \xrightarrow{T_\mu^{**}} & L_1(G)^{**} \\ \Gamma_E \uparrow & & \uparrow \Gamma_E \\ M(G) & \xrightarrow{M_\mu} & M(G) \end{array}$$

Indeed, given $\nu \in M(G)$ and $h \in L_\infty(G)$, we have

$$\begin{aligned} \langle \Gamma_E M_\mu(\nu), h \rangle &= \langle \Gamma_E(\mu \star \nu), h \rangle \\ &= \langle T_{\mu \star \nu}^{**} E, h \rangle \\ &= \langle E, T_{\mu \star \nu}^* h \rangle \\ &= \langle E, T_\nu^*(T_\mu^* h) \rangle \\ &= \langle T_\nu^{**} E, T_\mu^* h \rangle \\ &= \langle \Gamma_E(\nu), T_\mu^* h \rangle \\ &= \langle T_\mu^{**} \Gamma_E(\nu), h \rangle. \end{aligned}$$

Thus

$$(5) \quad \Gamma_E M_\mu = T_\mu^{**} \Gamma_E.$$

Now T_μ tauberian implies T_μ^{**} tauberian [9, Theorem 4.4.2]. Therefore $T_\mu^{**} \Gamma_E = \Gamma_E M_\mu$ is tauberian, and hence M_μ is tauberian, in both cases by [9, Proposition 2.1.3].

Similarly, denoting by $J : L_1(G) \rightarrow M(G)$ the natural isomorphic embedding, we have $JT_\mu = M_\mu J$. Hence, by [9, Proposition 2.1.3], if M_μ is tauberian, so is T_μ . \square

Corollary 3.8. *Let G be a LCA group. If the convolution operators T_μ is cotauberian, then M_μ is Fredholm.*

Proof. Let T_μ be cotauberian, then T_μ is Fredholm (Theorem 3.6). Therefore T_μ^{**} is Fredholm and $T_\mu^{**} \Gamma_E$ is upper semi-Fredholm. Then, by formula (5), M_μ is upper semi-Fredholm. Since $\mu = \lambda * \nu$ where $\lambda * \lambda = \lambda$ and ν is invertible [12, Theorem 1], we have that $M(G) = R(M_\mu) \oplus N(M_\mu)$. Hence M_μ is Fredholm. \square

Recall that an operator $T : L_1(G) \rightarrow L_1(G)$ is tauberian if and only if $m \in L_1(G)^{**}$ and $T^{**}m \in L_1(G)$ imply $m \in L_1(G)$.

Observation 3.9. *It was asked in [7] whether a convolution operator T_μ acting on $L_1(G)$ is tauberian when the measure μ satisfies the following condition:*

$$(6) \quad \nu \in M(G), \mu \star \nu \in L_1(G) \Rightarrow \nu \in L_1(G).$$

Next we will show that the answer to this question is negative.

Indeed, it was proved in [3] that there exists an atomic measure $\mu_0 \in M(\mathbb{T})$ such that T_{μ_0} is an injective non-tauberian operator, where \mathbb{T} denotes the unit circle. It is enough to choose μ_0 such that its Fourier-Stieltjes transform $\widehat{\mu}_0$ satisfies $0 \in \widehat{\mu_0(\mathbb{Z})} \setminus \widehat{\mu_0}(\mathbb{Z})$. The following argument, due to Doss [5], shows that μ_0 satisfies condition (6):

Every $\nu \in M(\mathbb{T})$ can be written as $\nu = \nu_1 + \nu_2$ with $\nu_1 \ll m$ and $\nu_2 \perp m$, where m is the Haar measure on \mathbb{T} . Since $\mu_0 \star \nu_1 \in L_1(\mathbb{T})$ and $\mu_0 \star \nu_2$ is supported in a m -null set, $T_{\mu_0} \nu \in L_1(\mathbb{T})$ if and only if $\nu_2 = 0$. Thus μ_0 satisfies (6).

It is not difficult to show that $\mu \star \tilde{f} = \tilde{\mu} \star f$ for $\mu \in M(G)$ and $f \in L_1(G)$. Also, for every normalized disjoint sequence $(f_n) \subset L_1(G)$, the sequence (\tilde{f}_n) is also normalized and disjoint. Therefore, it follows from [9, Theorem 4.1.3] that T_μ is tauberian if and only if so is $T_{\tilde{\mu}}$. Hence, by Theorem 3.7, the same happens for M_μ and $M_{\tilde{\mu}}$, and we get the following result:

Proposition 3.10. *Let G be a non-compact LCA group. Then*

- (i) $L_\mu : L_\infty(G) \rightarrow L_\infty(G)$ is tauberian if and only if it is cotauberian, and this is equivalent to L_μ invertible;
- (ii) $M_\mu : M(G) \rightarrow M(G)$ is tauberian if and only if it is invertible;
- (iii) $S_\mu : C_0(G) \rightarrow C_0(G)$ is cotauberian if and only if it is invertible.

Proof. (i) Note that $L_\mu = T_\mu^*$ and $T_{\tilde{\mu}}$ is tauberian if and only if so is its second conjugate $T_{\tilde{\mu}}^{**}$ [9, Theorem 4.4.2]. So the result follows from Theorem 2.1 and Corollary 3.5.

(ii) It is a consequence of Theorems 3.7 and 2.1, and Theorem 1.1.3 in [15].

(iii) Since $S_\mu^* = M_{\tilde{\mu}}$, the result follows from (ii). \square

Acknowledgements.

We are indebted to the referee for several comments that improved the paper.

REFERENCES

- [1] P. Aiena. *Fredholm and local spectral theory, with applications to multipliers*. Kluwer Academic Press, 2004.
- [2] R. Arens. *The adjoint of a bilinear operation*. Proc. Amer. Math. Soc. 2 (1951), 839–848.
- [3] L. Cely, E. M. Galego and M. González. *Tauberian convolution operators acting on $L_1(G)$* . J. Math. Anal. Appl. 446 (2017), 299–306.
- [4] C-H. Chu. *Matrix convolution operators on groups*. Lecture Notes in Math. 1956. Springer, 2008.
- [5] R. Doss. *Convolution of singular measures*. Studia Math. 45 (1973), 111–117.
- [6] F. Ghahramani, A. T. Lau and V. Losert. *Isometric isomorphisms between Banach algebras related to locally compact groups*. Trans. Amer. Math. Soc. 321 (1990), 273–283.
- [7] M. González. *Tauberian operators. Properties, applications and open problems*. In “Concrete operators, spectral theory, operators in harmonic analysis and approximation”, pp. 231–242. Operator Theory: Advances & Appl. 236. Birkhäuser, 2014.
- [8] M. González and A. Martínez-Abejón. *Tauberian operators on $L_1(\mu)$ spaces*. Studia Math. 125 (1997), 289–303.
- [9] M. González and A. Martínez-Abejón. *Tauberian operators*. Operator Theory: Advances and applications 194. Birkhäuser, 2010.
- [10] M. González and V. M. Onieva. *Characterizations of tauberian operators and other semigroups of operators*. Proc. Amer. Math. Soc. 108 (1990), 399–405.
- [11] M. González, E. Saksman and H.-O. Tylli. *Representing non-weakly compact operators*. Studia Math. 113 (1995), 265–282.
- [12] B. Host and F. Parreau. *Sur un problème de I. Glicksberg: Les idéaux fermés de type fini de $M(G)$* . Ann. Inst. Fourier (Grenoble) 28 (1978), 143–164.

- [13] W. B. Johnson, A. B. Nasser, G. Schechtman and T. Tkocz. *Injective tauberian operators on L_1 and operators with dense range on ℓ_∞* . Canad. Math. Bull. 58 (2015), 276–280.
- [14] N. Kalton, A. Wilansky. *Tauberian operators on Banach spaces*. Proc. Amer. Math. Soc. 57 (1976), 251–255.
- [15] R. Larsen. *An introduction to the theory of multipliers*. Springer, 1971.
- [16] A. T. Lau and V. Losert. *On the second conjugate algebra of $L_1(G)$ of a locally compact group*. J. London Math. Soc. 37 (1988), 464–470.
- [17] V. Losert, M. Neufang, J. Pachl and J. Steprāns. *Proof of the Ghahramani-Lau conjecture*. Advances in Math. 290 (2016), 709–738.
- [18] W. Rudin. *Fourier analysis on groups*. Wiley & Sons, 1962.
- [19] D. G. Tacon. *Generalized semi-Fredholm transformations*. J. Austral. Math. Soc. 347 (1983), 60–70.
- [20] S. Watanabe. *A Banach algebra which is an ideal in the second dual space*. Sci. Rep. Niigata Univ. Ser. A 11 (1974), 95–101.

DEPARTAMENTO DE MATEMÁTICA, INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, 05508-090 SÃO PAULO SP, BRAZIL
E-mail address: mlcelyp@unial.edu.co

DEPARTAMENTO DE MATEMÁTICA, INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, 05508-090 SÃO PAULO SP, BRAZIL
E-mail address: eloi@ime.usp.br

DEPARTAMENTO DE MATEMÁTICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD DE CANTABRIA, E-39071 SANTANDER, SPAIN
E-mail address: manuel.gonzalez@unican.es